

# CLOMAX ORAL



Active Ingredients: 1g/L Abamectin, 50g/L Closantel (as sodium salt) with 0.5g/L Selenium & 2.2g/L Cobalt.

## ADMINISTRATION

By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

VETMED Clomax Oral Drench is ready to use through standard drenching equipment. Rinse gun with water immediately after use.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching technique should be used. Check accuracy of drench guns regularly.

DOSE RATE: 1 mL/5 kg body weight delivering 0.2 mg/kg Abamectin, 10 mg/kg Closantel, 0.1 mg/kg Selenium and 0.44 mg/kg Cobalt.

Do not use in lambs under 15 kg or 4 weeks of age.

LIVEWEIGHT	DOSE VOLUME
15 kg	3 mL
16 – 20 kg	4 mL
21 – 25 kg	5 mL
26 – 30 kg	6 mL
31 – 35 kg	7 mL
36 – 40 kg	8 mL
41 – 45 kg	9 mL
46 – 50 kg	10 mL
51 – 55 kg	11 mL
56 – 60 kg	12 mL
61 – 65 kg	13 mL
66 – 70 kg	14 mL
71 – 75 kg	15 mL
Over 75 kg	add 1 mL / 5 kg

## WITHHOLD PERIODS

MEAT: 56 days, MILK: 56 days.

## STORAGE

Store at or below 25°C. Protect from light. Stability in cold temperatures has not been assessed. Discard unused product.

## EFFICACY

VETMED Clomax Oral Contains 1g/L Abamectin a broad spectrum drench of the macrocyclic lactone group, 50g/L Closantel (as sodium salt) a narrow spectrum salicylanilide drench and 0.5g/L Selenium and 2.2 g/L Cobalt. It has persistent activity up to 42 days against Barber's Pole (*Haemonchus contortus*).

At recommended doses, this product is effective against sensitive strains of the following mature and immature roundworms and liver fluke in sheep:

Barber's Pole (*Haemonchus contortus*), Stomach hair worm (*Trichostrongylus axei*), Black scour worm (*Trichostrongylus colubriformis* and *Trichostrongylus vitrinis*), Brown stomach worm (*Teladorsagia circumcincta* and *Teladorsagia trifurcata*), Small intestinal worm (*Cooperia oncophora*, *Cooperia surnabada* and *Cooperia punctata*), Thin-necked intestinal worm (*Nematodirus spathiger*), Liver fluke (*Fasciola hepatica*), Lungworm (*Dictyocaulus filaria*).

## PACK SIZES

VETMED Clomax Oral is available in 20L, 10L, 5L, 2.5L and 1L packs.

VETMED Clomax Oral is registered pursuant to the ACVM Act 1997, No. A011291. See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR100758. See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.



09 274 8848 | [www.vetmed.co.nz](http://www.vetmed.co.nz)

Registered to: Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland

+2

ABAMECTIN  
CLOSANTEL

BENEFITS

- Persistent activity for up to 42 days against Barber's Pole (*Haemonchus Contortus*)
- Oral drench for the treatment and control of roundworms and liver fluke in sheep
- Broad Spectrum – Oral Drench
- Dual Active – Roundworm and Liver Fluke control
- Developed in New Zealand for New Zealand



# CLOMAX ORAL



SHEEP

**MINERALISED ORAL DRENCH FOR THE CONTROL AND TREATMENT OF INTERNAL PARASITES, INCLUDING PERSISTENT ACTIVITY AGAINST BARBER'S POLE (*HAEMONCHUS CONTORTUS*), AND LIVER FLUKE (*FASCIOLA HEPATICA*) IN SHEEP.**

Active Ingredients: 1g/L Abamectin, 50g/L Closantel (as sodium salt) with 0.5g/L Selenium & 2.2g/L Cobalt.