

ABAMECTIN

POUR-ON FOR CATTLE

Active Ingredient: 10g/L Abamectin



ADMINISTRATION

Ready for use with standard Pour-On equipment. Check accuracy of Pour-On applicator regularly. A representative sample of animals should be weighed before treatment either with scales or weighband.

Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Apply along the backline from shoulders to tail.

This product should not be used on cattle which are under 100kg in weight or less than 16 weeks old.

DOSE: 1mL per 20kg bodyweight (500 micrograms Abamectin per kg bodyweight).

WEIGHT RANGE	DOSE VOLUME	WEIGHT RANGE	DOSE VOLUME
101 – 150 kg	7.5 mL	301 – 350 kg	17.5 mL
151 – 200 kg	10 mL	351 – 400 kg	20 mL
201 – 250 kg	12.5 mL	401 – 450 kg	22.5 mL
251 – 300 kg	15 mL	451 – 500 kg	25 mL

Cattle in excess of 500kg to be dosed at 1mL per 20kg bodyweight.

COMBINED USE

Combined use of drench actives may improve efficacy and slow the development of resistance. VETMED Abamectin Pour-On can be combined in equal volumes with VETMED Levamisole Pour-On (A10760) to produce a stable mixture for Pour-On dosing of cattle. The mixture contains 5.0g/L Abamectin and 100g/L Levamisole. This combined solution must be well mixed before use and dosed at a rate of 1mL/10kg bodyweight. The mixture remains stable for up to 3 months if stored as per label conditions.

RAIN

Treatment of animals with wet hides or when rain threatens should be avoided.

EFFICACY

VETMED Abamectin Pour-On for Cattle when used at the recommended dose level of 500 mg Abamectin per kg liveweight effectively controls the following parasites in cattle:

- Gastro-intestinal roundworms: adult and immature *Ostertagia* spp (including inhibited immatures), *Haemonchus* spp, *Trichostrongylus* spp, *Cooperia* spp, *Bunostomum* spp, *Oesophagostomum* spp, and adults of *Chabertia* spp, *Nematodirus* spp, *Trichuris* spp and *Strongyloides* spp.
- Lungworms: adult and immature *Dictyocaulus viviparus*.
- Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

WITHHOLD PERIODS

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within 35 days of the last treatment.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment.

STORAGE

Store in the original container, tightly closed and locked up in a safe place. Store below 25°C away from direct sunlight.

PACK SIZES

VETMED Abamectin Pour-On is available in 20L, 5L and 2.5L packs.

VETMED Abamectin Pour-On is registered pursuant to the ACVM Act 1997 No. A10786.

***INCREASED MILK PRODUCTION** - Increases Milk production in grazing dairy cows - Data has shown that milk production increases when cows or heifers infected with gastrointestinal nematode parasites are treated with an effective anthelmintic.



09 274 8848 | www.vetmed.co.nz


Registered to; Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland

+1

ABAMECTIN

BENEFITS

- Improves milk production*
- Convenient Pour-On application
- Broad spectrum activity
- Effective against sucking and biting lice (*Damalinia bovis*, *Linognathus vituli*)
- Effective against lungworm (adult and immaturely *Dictyocaulus viviparus*)
- Solvent base delivering high blood levels of Abamectin
- Dramatic visible results
- Developed in New Zealand for New Zealand



CATTLE

FOR THE TREATMENT AND CONTROL
OF INTERNAL AND EXTERNAL PARASITES
OF CATTLE. CONVENIENT POUR ON
APPLICATION. FAST ACTING - BROAD
SPECTRUM.

Active Ingredient: 10g/L Abamectin