

FLUMETHRIN

POUR-ON TICKICIDE FOR CATTLE AND DEER



Active Ingredient: 10g/L Flumethrin.

ADMINISTRATION

Apply VETMED Flumethrin Pour-On evenly along the midline of the back from the front of the shoulders to the tail head.

Use a plastic measuring cup or automatic drenching gun to apply.

Retreatment should be undertaken when ticks start to reappear.

For optimum control of deer, breeding hinds should be treated every three weeks from August to fawning. This program will reduce the number of nymphs reaching the adult stage, which infest fawn during November/December. Treatment of hinds can be continued after fawning if this is practical. Other stock should be treated every three weeks from August, or as required.

DOSE	
CATTLE (Liveweight)	
WEIGHT RANGE	DOSE VOLUME
Up to 200 kg	20 mL
200 – 300 kg	30 mL
301 – 400 kg	40 mL
Over 400 kg	50 mL

DEER	
WEIGHT RANGE	DOSE VOLUME
Up to 6 months	10 mL
Yearlings and adults	20 mL

WITHHOLD PERIODS

NO WITHHOLDING PERIOD for meat and milk from treated animals.

RAIN

Control of ticks is not affected by rain falling within a few hours of treatment.

STORAGE

Store below 25°C (room temperature). Do not freeze.

ADVERSE EFFECTS

Signs of mild irritation may be observed for 24 to 48 hours after treatment. Hair loss at the site of application may occur on deer which are shedding their winter coat at the time of treatment. This effect is temporary and can be minimised by spreading the dose over the entire backline. Growth of summer coat is not affected.

PACK SIZES

VETMED Flumethrin Pour-On is available in 10L herd, 5L and 1L packs.

VETMED Flumethrin Pour-On is registered pursuant to the ACVM Act 1997 No. A10816.




+1

FLUMETHRIN

BENEFITS

- Cost effective tickicide treatment
- Treats and controls all 3 stages of lifecycle
- Rainfast – not affected by rain falling within a few hours
- Easy to use Pour-On total body coverage
- Nil meat and milk withholding
- Persistent activity against ticks
- Dramatic visible results
- Developed in New Zealand for New Zealand



FLUMETHRIN

POUR-ON TICKICIDE FOR CATTLE AND DEER



CATTLE



DEER

SCIENTIFICALLY FORMULATED TO KILL TICKS. FOR THE CONTROL OF TICKS ON CATTLE AND AS AN AID IN THE TREATMENT OF TICKS ON DEER.

Active Ingredient: 10g/L Flumethrin.