

MOXIDECTIN POUR-ON



FAST ACTING - LONG LASTING - BROAD SPECTRUM

Active Ingredient: 5mg/mL Moxidectin.

ADMINISTRATION

By law the user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

SHAKE WELL BEFORE USE.

Invert drum three times before use. Administer 1mL per 10kg liveweight using any standard drenching equipment. Correct drenching technique should be used. Equipment should be checked for accuracy of dose delivery before drenching commences. Check accuracy of drench guns regularly.

DOSAGE RATE: 1mL per 10kg liveweight. This will administer the correct dose of 0.5mg Moxidectin/kg liveweight. The following table may be used as a guide for the dose volume to use.

LIVE WEIGHT	DOSE VOLUME	DOSES PER PACK		
		500mL	1L	5L
50 kg	5 mL	100	200	1000
60 kg	6 mL	83	166	833
70 kg	7 mL	71	142	714
80 kg	8 mL	62	125	625
90 kg	9 mL	55	111	555
100 kg	10 mL	50	100	500
110 kg	11 mL	45	90	454
120 kg	12 mL	41	83	416
130 kg	12 mL	38	76	384
140 kg	14 mL	35	71	357
150 kg	15 mL	33	66	333
200 kg	20 mL	25	50	250
250 kg	25 mL	20	40	200
300 kg	30 mL	16	33	166
350 kg	35 mL	14	28	142
400 kg	40 mL	12	25	125
450 kg	45 mL	11	22	111
500 kg	50 mL	10	20	100
550 kg	55 mL	9	18	90
600 kg	60 mL	8	16	83
650 kg	65 mL	7	15	76
700 kg	70 mL	7	14	71
750 kg	75 mL	6	13	66

FOR ANIMALS HEAVIER THAN 750KG, USE A VOLUME OF 1ML PER 10KG LIVELWEIGHT

WITHHOLD PERIODS

MILK: NIL, MEAT: Cattle - NIL, Deer - NIL, Bobby Calf - NIL.

STORAGE

Store below 25°C. Protect from Light.

RAIN

The efficacy of VETMED Moxidectin Pour-On is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However treatment of animals under these circumstances is not recommended practice.

PERSISTANT ACTIVITY

VETMED Moxidectin Pour-On, following a single application at the recommended dose rate prevents reinfection with the following parasite species:

CATTLE		DEER	
Parasite Species	Persistent protection period	Parasite Species	Persistent protection period
<i>Dictyocaulus viviparus</i>	42 days	<i>Dictyocaulus ekerti</i>	42 days
<i>Ostertagia ostertagi (O.lyrata)</i>	35 days	<i>Ostertagia-like spp*</i>	42 days
<i>Haemonchus spp</i>	28 days	<i>Oesophagostomum venulosum</i>	42 days
<i>Trichostrongylus axei</i>	28 days	<i>Chabertia ovina</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days		
<i>Bunostomum phlebotomum</i>	42 days		

**Ostertagia leptospicularis* and *Ostertagia*-like species including *Spiculopteragia assymetrica*, *Spiculopteragia spiculoptera* and *Skrjabinagia kolchida*.

VETMED Moxidectin Pour-On is registered pursuant to the ACVM Act 1997 No. A10783. ERMA Approval No. HSR001850.

PACK SIZES

VETMED Moxidectin Pour-On is available in 20L, 5L, 1L and 500mL packs.




09 274 8848 | www.vetmed.co.nz

Registered to: Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland

+1 MOXIDECTIN

BENEFITS

- Contains Moxidectin – a potent macrocyclic lactone (ML)
- Improves milk production
- Rainfast
- Persistent activity
- Non-Irritant formulation
- Broad spectrum
- Safe to use concurrently with other treatments
- Developed in New Zealand for New Zealand



MOXIDECTIN POUR-ON

FAST ACTING - LONG LASTING - BROAD SPECTRUM



CATTLE



DEER

FOR THE TREATMENT AND CONTROL OF INTERNAL AND EXTERNAL PARASITES OF CATTLE (INCLUDING LACTATING DAIRY CATTLE) AND FOR THE TREATMENT AND CONTROL OF LUNGWORM AND ROUNDWORMS OF DEER.

Active Ingredient: 5mg/mL Moxidectin.